

ORGANIC CHEMISTRY

Article

Received: 2 June 2026 | Revised: 14 August 2026 |

Accepted: 18 August 2026 | Published online: 28 August 2026

UDC 544.42+519.242.7

<https://doi.org/10.31489/2959-0663/3-26-7>

Elena V. Salimova , Rinat R. Gubaidullin , Lyudmila V. Parfenova* 

Institute of Petrochemistry and Catalysis, Ufa Federal Research Center of the Russian Academy of Sciences, Ufa, Russia

(*Corresponding author's e-mail: luda_parfenova@mail.ru)

Synthesis of Amphiphilic Derivatives of Fusidane Triterpenoids

Fusidic acid exhibits bacteriostatic activity and can exert bactericidal effects at high concentrations, primarily against Gram-positive bacteria, including methicillin-resistant *Staphylococcus aureus*. This provides opportunities for functionalization of its reactive sites with pharmacophoric groups to develop semisynthetic analogs with potential antibacterial activity. The development of amphiphilic fusidic acid derivatives with improved bioavailability and biological activity is therefore of considerable interest. In this work, amphiphilic fusidic acid (FA) derivatives were synthesized via quaternization of tertiary amines and pyridines (triethylamine, pyridine, N,N-dimethylaminopyridine, and N-methylpiperidine) with bromoethyl fusidate and 3-bromopropylamine methylfusidate. The carboxylic group of FA was converted into its bromoethyl ester using 1,2-dibromoethane and K₂CO₃ in dry acetone under reflux conditions. Reaction of the resulting bromoethyl fusidate with tertiary amines and pyridines in DMF at 70 °C afforded the corresponding ammonium salts in 62–70% yields. The FA bromoamine, obtained by reductive amination, was subsequently reacted with N-containing compounds in DMF under heating to afford the target products in 60–67% yields. The structures of the novel FA analogs were confirmed by NMR spectroscopy and mass spectrometry. The obtained amphiphilic FA analogs may provide a basis for further development of antimicrobial agents.

Keywords: triterpenoids, fusidic acid, amphiphilic derivatives, tertiary amines, quaternization reaction, quaternary ammonium salts, reductive amination.

Introduction

For many years, amphiphilic quaternary ammonium compounds have been widely recognized for their biological activity, primarily as antimicrobial agents [1–11]. Additionally, this class of compounds exhibits antiviral, anticancer [12–17], and membrane-tropic properties [18–20]. Evidently, one of the key reasons underlying the biological activity of amphiphilic compounds is the presence of pharmacophoric ammonium groups, and in some cases, hydrophobic polyalkyl fragments. In this context, functionalizing natural polycyclic terpenoids which possess a hydrophobic carbocyclic core and inherent biological activity—with ammonium groups is of considerable interest.

Fusidane triterpenoids represent a small group of natural compounds produced by various fungal species, each displaying antibacterial activity to varying degrees [21–23]. Among them, fusidic acid (FA) holds a special place, having found clinical application in the therapy of staphylococcal infections. Since its discovery, multiple functionalizations of FA's reactive centers with pharmacophore groups have been performed, yielding analogs with diverse biological activities [24–29]. However, to the best of our knowledge amphiphilic derivatives of fusidane triterpenoids have not been reported. Taking this into account, within our ongoing research on synthetic transformations of FA [28–34], we have synthesized novel amphiphilic analogs via quaternization of tertiary aliphatic amines and pyridines with bromoethyl fusidate and 3-bromopropylamine methylfusidate.

Experimental

^1H and ^{13}C NMR spectra were recorded on a Bruker Avance II 500 HD Ascend spectrometer (Germany) [500.17 MHz (^1H), 125.78 MHz (^{13}C)]. Samples were prepared in standard 5 mm diameter ampoules. One- and two-dimensional NMR spectra (COSY, HSQC, HMBC, NOESY) were acquired using standard pulse sequences. Melting points were determined on a Stuart SMP3 apparatus (Germany). Optical rotations were measured on a Perkin-Elmer 341 polarimeter (Germany). MALDI TOF/TOF mass spectra were obtained on a Bruker AutoflexTM III Smartbeam spectrometer (Germany) using 3-(4-hydroxy-3,5-dimethoxyphenyl)prop-2-enoic acid (sinapinic acid) as matrix. TLC was performed on Sorbfil plates (Sorbpolymer Ltd., Russia) using $\text{CHCl}_3/\text{MeOH}$ (40:1); spots were visualized with 10 % H_2SO_4 followed by heating at 100–120 °C for 2–3 min. Silica gel (50–160 μm , Sorbpolymer Ltd., Russia) was used for column chromatography. Commercially available reagents of “chemically pure” or “analytical grade” (Acros Organics) were employed. Fusidic acid was purchased from Hangzhou Hyper Chemicals Limited (purity 99.3 %). Methyl fusidate (**7**) and its dioxo derivative (**8**) were prepared according to a literature procedure [30].

2-Bromoethyl (Z)-2-((3S,4S,8S,10S,11R,14S,16S)-16-acetoxy-3,11-dihydroxy-4,8,10,14-tetramethylhexadecahydro-17H-cyclopenta[a]phenanthren-17-ylidene)-6-methylhept-5-enoate (2). K_2CO_3 (0.04 g, 0.28 mmol) and 1,2-dibromoethane (0.17 mL, 1.95 mmol) were added to a solution of fusidic acid **1** (0.20 g, 0.40 mmol) in 20 mL of dry acetone. The reaction mixture was heated at 56 °C for 8 h. Acetone was removed under reduced pressure, the residue was dissolved in 30 mL of EtOAc, washed with H_2O and dried over CaCl_2 . The solvent was evaporated under reduced pressure and the residue was put under vacuum. Yield of compound **2**: 0.23 g (97 %), white powder. Mp 105–107 °C, $[\alpha]_{\text{D}}^{20} = -17^\circ$ (c 0.82, CHCl_3). Spectral data are given in Ref. [35].

Compounds 3–6 (general procedure). Bromoethyl ester **2** (1.00 g, 1.67 mmol) was dissolved in 5 mL of DMF and the corresponding tertiary amine (12 mmol) was added: triethylamine (1.21 g), pyridine (0.95 g), *N,N*-dimethylaminopyridine (1.47 g), or *N*-methylpiperidine (1.19 g). The reaction mixture was stirred at 70 °C for 8 h. After completion, the mixture was purified by column chromatography on SiO_2 , eluting with $\text{CHCl}_3/\text{MeOH}$ (10:1).

2-(((Z)-2-((3S,4S,8S,10S,11R,14S,16S)-16-acetoxy-3,11-dihydroxy-4,8,10,14-tetramethylhexadecahydro-17H-cyclopenta[a]phenanthren-17-ylidene)-6-methylhept-5-enoyl)oxy)-*N,N,N*-triethylethan-1-aminium bromide (3)

Yield 0.72 g (62 %), white powder, mp: 187–189 °C, $[\alpha]_{\text{D}}^{19} -10^\circ$ (c 0.86, MeOH). ^1H NMR (MeOD): $\delta = 0.85$ s (3H, H^{18}), 0.88 d (3H, H^{28} , J 6.3 Hz), 0.91 s (3H, H^{19}), 0.97–1.09 m (2H, H^{6a} , H^{7a}), 1.12–1.22 m (1H, H^{15a}), 1.32 s (3H, H^{30}), 1.37 t (9H, H^4 , H^6 , H^8 , J 7.5 Hz), 1.38–1.47 m (1H, H^{1a}), 1.41–1.51 m (1H, H^5), 1.46–1.55 m (1H, H^9), 1.53–1.61 m (1H, H^{6b}), 1.55 s (3H, H^{27}), 1.61 s (3H, H^{26}), 1.61–1.69 m (1H, H^{7b}), 1.62–1.81 m (2H, H^2), 1.71–1.82 m (1H, H^{12a}), 1.87–2.21 m (2H, H^{23}), 1.96 s [3H, $\text{O}-\text{C}(\text{O})\text{CH}_3$], 2.01–2.12 m (1H, H^4), 2.05–2.16 m (1H, H^{15b}), 2.06–2.21 m (1H, H^{1b}), 2.24–2.35 m (1H, H^{12b}), 2.25–2.43 m (2H, H^{22}), 3.03 d (1H, H^{13} , J 10.1 Hz), 3.38–3.63 m (6H, $\text{H}^{3'}$, H^5 , H^7), 3.62–3.72 m (1H, H^3), 3.65–3.81 m (2H, H^2), 4.15–4.26 m (1H, H^{1a}), 4.25–4.36 m (1H, H^{11}), 4.50–4.61 m (1H, H^{1b}), 5.01 t (1H, H^{24} , J 6.4 Hz), 5.77 d (1H, H^{16} , J 6.4 Hz). ^{13}C NMR (MeOD): $\delta = 8.2$ (C^4 , C^6 , C^8), 16.1 (C^{28}), 17.6 (C^{18}), 18.0 (C^{27}), 21.0 (C^6), 21.3 [$\text{O}-\text{C}(\text{O})\text{CH}_3$], 23.5 (C^{19}), 23.6 (C^{30}), 25.7 (C^{26}), 28.8 (C^{23}), 28.9 (C^{22}), 29.8 (C^2), 29.9 (C^1), 31.5 (C^7), 35.4 (C^4), 35.5 (C^{12}), 36.6 (C^{10}), 36.7 (C^5), 39.0 (C^{15}), 39.5 (C^8), 44.7 (C^{13}), 48.7 (C^{14}), 49.3 (C^9), 54.4 ($\text{C}^{3'}$, C^5 , C^7), 55.7 (C^{21}), 57.5 (C^{11}), 67.8 (C^{11}), 71.2 (C^3), 74.0 (C^{16}), 122.9 (C^{24}), 128.7 (C^{20}), 132.7 (C^{25}), 151.3 (C^{17}), 168.7 (C^{21}), 170.9 [$\text{O}-\text{C}(\text{O})\text{CH}_3$].

Calcd, %: C 64.62; H 9.18; Br 11.02; N 1.93; O 13.24. $\text{C}_{39}\text{H}_{66}\text{BrNO}_6$. Found, %: C 64.56; H 9.20; Br 11.05; N 1.92. m/z 747.4 [$\text{M}+\text{Na}$]⁺, calcd for [$\text{C}_{39}\text{H}_{66}\text{BrNO}_6\text{Na}$]⁺ 747.4.

1-(2-(((Z)-2-((3S,4S,8S,10S,11R,14S,16S)-16-acetoxy-3,11-dihydroxy-4,8,10,14-tetramethylhexadecahydro-17H-cyclopenta[a]phenanthren-17-ylidene)-6-methylhept-5-enoyl)oxy)ethyl)pyridin-1-ium bromide (4)

Yield 0.77 g (68 %), white powder, mp: 179–181 °C, $[\alpha]_{\text{D}}^{19} +9^\circ$ (c 0.95, CHCl_3). ^1H NMR (CDCl_3): $\delta = 0.87$ s (3H, H^{18}), 0.88 d (3H, H^{28} , J 6.5 Hz), 0.97 s (3H, H^{19}), 1.03–1.11 m (1H, H^{7a}), 1.05–1.13 m (1H, H^{6a}), 1.14–1.23 m (1H, H^{15a}), 1.36 s (3H, H^{30}), 1.43–1.50 m (1H, H^{1a}), 1.47–1.55 m (1H, H^5), 1.52 s (3H, H^{27}), 1.53–1.60 m (1H, H^9), 1.57–1.67 m (1H, H^{6b}), 1.62–1.70 m (1H, H^{7b}), 1.63 s (3H, H^{26}), 1.66–1.89 m (2H, H^2), 1.75–1.84 m (1H, H^{12a}), 1.94 s [3H, $\text{O}-\text{C}(\text{O})\text{CH}_3$], 2.01–2.16 m (2H, H^{23}), 2.04–2.13 m (1H, H^4), 2.06–2.17 m (1H, H^{15b}), 2.07–2.17 m (1H, H^{1b}), 2.19–2.31 m (1H, H^{12b}), 2.28–2.44 m (2H, H^{22}), 3.05 d (1H, H^{13} , J 8.5 Hz), 3.63–3.69 m (1H, H^3), 4.25–4.35 m (1H, H^{11}), 4.36–4.49 m (1H, H^{1a}), 4.61–4.71 m (1H, H^{1b}), 4.95 t (1H, H^{24} ,

J 7.1 Hz), 5.03-5.18 m (2H, H²), 5.66 d (1H, H¹⁶, J 8.5 Hz), 8.19 t (2H, H^{4'}, H^{6'}, J 7.5 Hz), 8.67 t (1H, H^{5'}, J 7.5 Hz), 9.14 d (2H, H^{3'}, H^{7'}, J 5.5 Hz). ¹³C NMR (CDCl₃), δ =15.7 (C²⁸), 17.3 (C²⁷), 17.5 (C¹⁸), 20.7 [O-C(O)CH₃], 21.1 (C⁶), 22.8 (C³⁰), 23.6 (C¹⁹), 25.7 (C²⁶), 28.4 (C²³), 28.7 (C²²), 29.6 (C¹, C²), 31.1 (C⁷), 35.2 (C⁴), 35.7 (C¹²), 36.4 (C¹⁰), 36.9 (C⁵), 38.9 (C¹⁵), 39.5 (C⁸), 44.5 (C¹³), 48.6 (C¹⁴), 49.4 (C⁹), 60.2 (C²), 62.9 (C¹), 67.5 (C¹¹), 71.2 (C³), 74.3 (C¹⁶), 122.7 (C²⁴), 128.6 (C^{4'}, C^{6'}), 128.7 (C²⁰), 132.6 (C²⁵), 145.2 (C^{3'}, C^{7'}), 146.4 (C⁵), 151.0 (C¹⁷), 169.0 (C²¹), 171.0 [O-C(O)CH₃]. Calcd, %: C 64.95; H 8.03; Br 11.37; N 1.99; O 13.66. C₃₈H₅₆BrNO₆. Found, %: C 64.87; H 8.05; Br 11.40; N 1.98. m/z 724.4 [M+Na]⁺, calcd for [C₃₈H₅₆BrNO₆Na]⁺ 724.3.

1-(2-(((Z)-2-((3S,4S,8S,10S,11R,14S,16S)-16-acetoxy-3,11-dihydroxy-4,8,10,14-tetramethylhexadecahydro-17H-cyclopenta[a]phenanthren-17-ylidene)-6-methylhept-5-enoyl)oxy)ethyl)-4-(dimethylamino)pyridin-1-ium bromide (5)

Yield 0.78 g (65 %), white powder, mp: 181-183°C, $[\alpha]_D^{19}$ +7° (c 0.89, CHCl₃). ¹H NMR (CDCl₃): δ =0.76 s (3H, H¹⁸), 0.80 d (3H, H²⁸, J 6.5 Hz), 0.83 s (3H, H¹⁹), 0.91-1.05 m (2H, H^{6a}, H^{7a}), 1.02-1.12 m (1H, H^{15a}), 1.25 s (3H, H³⁰), 1.30-1.39 m (1H, H^{1a}), 1.33-1.43 m (1H, H⁵), 1.38-1.45 m (1H, H⁹), 1.40 s (3H, H²⁷), 1.44-1.55 m (1H, H^{6b}), 1.50-1.64 m (1H, H^{7b}), 1.51 s (3H, H²⁶), 1.60-1.77 m (2H, H²), 1.62-1.72 m (1H, H^{12a}), 1.83 s [3H, O-C(O)CH₃], 1.92-2.02 m (2H, H²³), 1.97-2.08 m (1H, H⁴), 1.99-2.07 m (1H, H^{15b}), 2.07-2.17 m (1H, H^{1b}), 2.18-2.30 m (2H, H²²), 2.19-2.29 m (1H, H^{12b}), 2.97 d (1H, H¹³, J 10.8 Hz), 3.18 s (6H, H⁸, H⁹), 3.54-3.63 m (1H, H³), 4.13-4.23 m (1H, H^{1a}), 4.19-4.27 m (1H, H¹¹), 4.36-4.46 m (1H, H^{1b}), 4.48-4.59 m (1H, H^{2a}), 4.64-4.75 m (1H, H^{2b}), 4.86 t (1H, H²⁴, J 6.8 Hz), 5.60 d (1H, H¹⁶, J 6.8 Hz), 6.88 d (2H, H^{4'}, H^{6'}, J 5.4 Hz), 8.39 d (2H, H^{3'}, H^{7'}, J 5.4 Hz). ¹³C NMR (CDCl₃), δ =16.0 (C²⁸), 17.5 (C²⁷), 17.8 (C¹⁸), 21.0 [O-C(O)CH₃], 21.1 (C⁶), 23.4 (C³⁰), 23.6 (C¹⁹), 25.7 (C²⁶), 28.7 (C²³), 28.9 (C²²), 29.8 (C²), 29.9 (C¹), 31.3 (C⁷), 35.2 (C⁴), 35.5 (C¹²), 36.5 (C¹⁰), 36.9 (C⁵), 38.9 (C¹⁵), 39.4 (C⁸), 40.4 (C^{8'}, C^{9'}), 44.4 (C¹³), 48.5 (C¹⁴), 49.4 (C⁹), 56.1 (C²), 63.2 (C¹), 67.5 (C¹¹), 70.9 (C³), 74.1 (C¹⁶), 108.1 (C^{4'}, C^{6'}), 123.1 (C²⁴), 128.7 (C²⁰), 132.2 (C²⁵), 142.7 (C^{3'}, C^{7'}), 150.4 (C¹⁷), 156.4 (C^{5'}), 169.1 (C²¹), 170.6 [O-C(O)CH₃]. Calcd, %: C 64.42; H 8.24; Br 10.71; N 3.76; O 12.87. C₄₀H₆₁BrN₂O₆. Found, %: C 64.49; H 8.22; Br 10.68; N 3.75. m/z 767.5 [M+Na]⁺, calcd for [C₄₀H₆₁BrN₂O₆Na]⁺ 767.4.

1-(2-(((Z)-2-((3S,4S,8S,10S,11R,14S,16S)-16-acetoxy-3,11-dihydroxy-4,8,10,14-tetramethylhexadecahydro-17H-cyclopenta[a]phenanthren-17-ylidene)-6-methylhept-5-enoyl)oxy)ethyl)-1-methylpiperidin-1-ium bromide (6)

Yield 0.81 g (70 %), white powder, mp: 173-175°C, $[\alpha]_D^{20}$ -12° (c 0.67, MeOH). ¹H NMR (MeOD): δ =0.88 s (3H, H¹⁸), 0.89 d (3H, H²⁸, J 6.4 Hz), 0.95 s (3H, H¹⁹), 1.04-1.14 m (2H, H^{6a}, H^{7a}), 1.21-1.28 m (1H, H^{15a}), 1.36 s (3H, H³⁰), 1.38-1.47 m (2H, H⁵), 1.44-1.51 m (1H, H^{1a}), 1.49-1.57 m (1H, H⁵), 1.52-1.63 m (4H, H^{4'}, H^{6'}), 1.53-1.58 m (1H, H⁹), 1.55-1.64 m (1H, H^{6b}), 1.58 s (3H, H²⁷), 1.66 s (3H, H²⁶), 1.66-1.72 m (1H, H^{7b}), 1.71-1.86 m (2H, H²), 1.76-1.85 m (1H, H^{12a}), 1.96 s [3H, O-C(O)CH₃], 1.97-2.19 m (2H, H²³), 2.05-2.14 m (1H, H⁴), 2.09-2.20 m (1H, H^{1b}), 2.11-2.20 m (1H, H^{15b}), 2.25-2.33 m (1H, H^{12b}), 2.35-2.50 m (2H, H²²), 2.44-2.53 m (4H, H^{3'}, H^{7'}), 2.64 t (2H, H^{2'}, J 6.1 Hz), 3.02 d (1H, H¹³, J 11.6 Hz), 3.44 s (3H, H⁸), 3.68-3.73 m (1H, H³), 4.03-4.09 m (1H, H^{1a}), 4.25-4.30 m (1H, H^{1b}), 4.30-4.35 m (1H, H¹¹), 5.06 t (1H, H²⁴, J 6.8 Hz), 5.81 d (1H, H¹⁶, J 7.8 Hz). ¹³C NMR (MeOD): δ =15.9 (C²⁸), 17.6 (C²⁷), 17.8 (C¹⁸), 20.9 (C⁶), 21.0 [O-C(O)CH₃], 23.3 (C¹⁹), 23.7 (C³⁰), 23.9 (C⁵), 25.5 (C^{4'}, C^{6'}), 25.7 (C²⁶), 28.4 (C²³), 28.9 (C²²), 29.8 (C²), 30.0 (C¹), 31.8 (C⁷), 35.5 (C¹²), 35.6 (C⁴), 36.6 (C⁵), 36.7 (C¹⁰), 38.9 (C¹⁵), 39.4 (C⁸), 44.0 (C¹³), 48.6 (C¹⁴), 49.3 (C⁹), 50.5 (C^{8'}), 54.6 (C^{3'}, C^{7'}), 56.8 (C²), 61.6 (C¹), 68.1 (C¹¹), 71.4 (C³), 74.4 (C¹⁶), 123.1 (C²⁴), 130.2 (C²⁰), 132.4 (C²⁵), 148.3 (C¹⁷), 170.0 (C²¹), 170.5 [O-C(O)CH₃]. Calcd, %: C 64.80; H 8.92; Br 11.05; N 1.94; O 13.28. C₃₉H₆₄BrNO₆. Found, %: C 64.74; H 8.94; Br 11.07; N 1.95. m/z 744.5 [M+Na]⁺, calcd for [C₃₉H₆₄BrNO₆Na]⁺ 744.4.

Methyl (Z)-2-((3S,4S,8S,10S,11R,14S,16S)-16-acetoxy-3-((3-bromopropyl)amino)-11-hydroxy-4,8,10,14-tetramethylhexadecahydro-17H-cyclopenta[a]phenanthren-17-ylidene)-6-methylhept-5-enoate (9). A mixture of diketone **8** (1.00 g, 1.92 mmol), titanium(IV) isopropoxide (0.23 g, 0.65 mmol) and 3-bromopropylamine (0.40 g, 2.89 mmol) in 10 mL of absolute methanol was stirred under an argon atmosphere at room temperature for 8 h. Sodium borohydride (0.15 g, 3.96 mmol) was then added at -78 °C and the resulting mixture was stirred for an additional 2 h while warming to 24 °C. Water (5 mL) was added to the reaction mixture, which was stirred at room temperature for 30 min. The precipitate was filtered off, the organic layer was concentrated under reduced pressure and placed under vacuum. The residue was purified by column chromatography on SiO₂, eluting with CHCl₃/MeOH (10:1).

Methyl (Z)-2-((3S,4S,8S,10S,11R,14S,16S)-16-acetoxy-3-((3-bromopropyl)amino)-11-hydroxy-4,8,10,14-tetramethylhexadecahydro-17H-cyclopenta[a]phenanthren-17-ylidene)-6-methylhept-5-enoate (9)

Yield 1.05 g (85 %), yellow powder, mp: 198-200°C, $[\alpha]_{\text{D}}^{19} +12^\circ$ (*c* 0.73, CHCl₃). ¹H NMR (CDCl₃): δ =0.88 s (3H, H¹⁸), 1.10 d (3H, H²⁸, *J* 6.8 Hz), 1.12 s (3H, H¹⁹), 1.14-1.27 m (1H, H^{7a}, 2H, H²), 1.15-1.26 m (1H, H^{15a}), 1.34 s (3H, H³⁰), 1.49-1.60 m (1H, H⁹), 1.58-1.66 m (1H, H^{7b}), 1.59 s (3H, H²⁷), 1.59-1.72 m (2H, H⁶), 1.64-1.71 m (1H, H^{1a}), 1.66 s (3H, H²⁶), 1.71-1.86 m (2H, H⁴, H^{12a}), 1.74-1.79 m (1H, H^{2a}), 1.87-1.97 m (1H, H^{1a}), 1.94-2.14 m (2H, H²³), 1.97 s [3H, O-C(O)CH₃], 2.01-2.11 m (3H, H^{1b}, H^{2b}, H⁵), 2.10-2.21 m (1H, H^{15b}), 2.21-2.29 m (1H, H^{12b}), 2.32-2.48 m (2H, H²²), 2.56-2.66 m (1H, H^{1b}), 2.66-2.74 m (1H, H^{3a}), 3.04 d (1H, H¹³, *J* 11.0 Hz), 3.04-3.15 m (1H, H³), 3.32-3.38 m (1H, H^{3b}), 3.63 s (3H, COOCH₃), 4.22-4.35 m (1H, H¹¹), 5.08 t (1H, H²⁴, *J* 7.5 Hz), 5.82 d (1H, H¹⁶, *J* 8.1 Hz). ¹³C NMR (CDCl₃): δ =14.2 (C²⁸), 17.1 (C¹⁸), 17.7 (C²⁷), 18.7 (C⁶), 20.9 [O-C(O)CH₃], 22.3 (C³⁰), 25.7 (C²⁶), 28.4 (C²³), 28.6 (C¹⁹), 28.8 (C²²), 29.7 (C⁷, C²), 29.9 (C²), 30.8 (C⁴), 31.6 (C¹), 36.3 (C¹²), 36.8 (C¹⁰), 38.7 (C³), 38.9 (C¹⁵), 39.9 (C⁸), 42.1 (C⁵), 43.8 (C¹³), 48.8 (C¹), 49.1 (C¹⁴), 50.5 (C⁹), 51.4 (COOCH₃), 66.7 (C¹¹), 67.4 (C³), 74.4 (C¹⁶), 123.0 (C²⁴), 130.3 (C²⁰), 132.6 (C²⁵), 147.8 (C¹⁷), 170.6 (C²¹), 171.0 [O-C(O)CH₃]. Calcd, %: C 64.60; H 8.67; Br 12.28; N 2.15; O 12.29. C₃₅H₅₆BrNO₅. Found, %: C 64.66; H 8.66; Br 12.25; N 2.16. *m/z* 672.4 [M+Na]⁺, calcd for [C₃₅H₅₆BrNO₅Na]⁺ 672.3.

Compounds 10–13 (general procedure). Bromoamine **9** (1.00 g, 1.54 mmol) was dissolved in 5 mL of DMF and the corresponding tertiary amine (11 mmol) was added: triethylamine (1.11 g), pyridine (0.87 g), N,N-dimethylaminopyridine (1.35 g), or N-methylpiperidine (1.09 g). The reaction mixture was stirred at 70 °C for 8 h. After completion, the mixture was purified by column chromatography on SiO₂, eluting with CHCl₃/MeOH (4:1).

3-(((3S,4S,8S,10S,11R,14S,16S, Z)-16-acetoxy-11-hydroxy-17-(1-methoxy-6-methyl-1-oxohept-5-en-2-ylidene)-4,8,10,14-tetramethylhexadecahydro-1H-cyclopenta[a]phenanthren-3-yl)amino)-N,N,N-triethylpropan-1-aminium bromide (10)

Yield 0.66 g (60 %), white powder, mp: 195-197°C, $[\alpha]_{\text{D}}^{19} +8^\circ$ (*c* 0.99, MeOH). ¹H NMR (MeOD): δ =0.88 s (3H, H¹⁸), 1.11 s (3H, H¹⁹), 1.17-1.29 m (1H, H^{15a}), 1.18 d (3H, H²⁸, *J* 6.5 Hz), 1.20-1.36 m (1H, H^{7a}), 1.35 s (3H, H³⁰), 1.44 t (9H, H⁵, H⁷, H⁹, *J* 7.2 Hz), 1.52-1.62 m (1H, H⁹), 1.55-1.64 m (2H, H⁶), 1.56-1.66 m (2H, H^{2a}, H^{7b}), 1.60 s (3H, H²⁷), 1.63-1.74 m (1H, H^{1a}), 1.67 s (3H, H²⁶), 1.72-1.86 s (2H, H²), 1.74-1.88 m (2H, H⁴, H^{12a}), 1.75-1.86 m (1H, H^{2b}), 1.92-2.04 m (1H, H^{1a}), 1.97-2.19 m (2H, H²³), 1.98 s [3H, O-C(O)CH₃], 2.04-2.14 m (1H, H^{1b}), 2.06-2.16 m (1H, H⁵), 2.10-2.23 m (1H, H^{15b}), 2.25-2.37 m (1H, H^{12b}), 2.36-2.53 m (2H, H²²), 2.63-2.74 m (1H, H^{1b}), 2.96-3.06 m (1H, H¹³), 3.18 t (6H, H⁴, H⁶, H⁸, *J* 6.1 Hz), 3.19-3.31 m (1H, H³), 3.64 s (3H, COOCH₃), 4.14-4.26 m (2H, H³), 4.32-4.47 m (1H, H¹¹), 5.11 t (1H, H²⁴, *J* 6.8 Hz), 5.83 d (1H, H¹⁶, *J* 8.1 Hz). ¹³C NMR (MeOD): δ =8.7 (C⁵, C⁷, C⁹), 14.8 (C²⁸), 17.1 (C¹⁸), 17.8 (C²⁷), 18.7 (C⁶), 21.0 [O-C(O)CH₃], 22.6 (C³⁰), 22.7 (C²), 25.8 (C²⁶), 28.4 (C²³), 28.7 (C¹⁹), 28.9 (C²²), 29.7 (C², C⁷), 31.1 (C⁴), 31.9 (C¹), 36.2 (C¹²), 36.8 (C¹⁰), 38.9 (C¹⁵), 39.8 (C⁸), 42.0 (C⁵), 43.9 (C¹³), 46.2 (C⁴, C⁶, C⁸), 48.5 (C¹⁴), 49.1 (C¹), 50.5 (C⁹), 51.4 (COOCH₃), 67.0 (C¹¹), 67.6 (C³), 68.1 (C³), 74.3 (C¹⁶), 123.1 (C²⁴), 130.3 (C²⁰), 132.5 (C²⁵), 147.9 (C¹⁷), 170.4 (C²¹), 170.8 [O-C(O)CH₃]. Calcd, %: C 65.49; H 9.52; Br 10.63; N 3.73; O 10.64. C₄₁H₇₁BrN₂O₅. Found, %: C 65.56; H 9.50; Br 10.61; N 3.72. *m/z* 773.5 [M+Na]⁺, calcd for [C₄₁H₇₁BrN₂O₅Na]⁺ 773.4.

1-(3-(((3S,4S,8S,10S,11R,14S,16S, Z)-16-acetoxy-11-hydroxy-17-(1-methoxy-6-methyl-1-oxohept-5-en-2-ylidene)-4,8,10,14-tetramethylhexadecahydro-1H-cyclopenta[a]phenanthren-3-yl)amino)propyl)pyridin-1-ium bromide (11)

Yield 0.75 g (67 %), white powder, mp: 182-184°C, $[\alpha]_{\text{D}}^{19} -11^\circ$ (*c* 1.12, MeOH). ¹H NMR (MeOD): δ =0.76 s (3H, H¹⁸), 1.01 d (3H, H²⁸, *J* 6.5 Hz), 1.02 s (3H, H¹⁹), 1.03-1.20 m (1H, H^{15a}), 1.09-1.19 m (1H, H^{7a}), 1.19 s (3H, H³⁰), 1.39-1.45 m (1H, H⁹), 1.45 s (3H, H²⁷), 1.46-1.59 m (2H, H^{1a}, H^{7b}), 1.53 s (3H, H²⁶), 1.55-1.70 m (2H, H⁶), 1.63-1.82 m (1H, H^{12a}), 1.69-1.76 m (1H, H^{2a}), 1.73-1.87 m (1H, H⁵), 1.76-1.91 m (1H, H^{1b}, 2H, H²), 1.83 s [3H, O-C(O)CH₃], 1.84-2.10 m (2H, H²³), 1.87-1.98 m (1H, H^{2b}), 1.89-2.10 m (1H, H^{1a}), 1.97-2.12 m (1H, H^{15b}), 2.12-2.29 m (1H, H^{12b}), 2.21-2.52 m (2H, H²²), 2.34-2.46 m (1H, H⁴), 2.37-2.53 m (1H, H^{1b}), 2.91 d (1H, H¹³, *J* 12.5 Hz), 3.05-3.15 m (1H, H³), 3.50 s (3H, COOCH₃), 4.14-4.17 m (2H, H³), 4.16-4.27 m (1H, H¹¹), 4.95 t (1H, H²⁴, *J* 6.5 Hz), 5.68 d (1H, H¹⁶, *J* 7.5 Hz), 8.03 t (2H, H⁵, H⁷, *J* 7.5 Hz), 8.43 t (1H, H⁶, *J* 7.5 Hz), 9.19 d (2H, H⁴, H⁸, *J* 5.3 Hz). ¹³C NMR (MeOD): δ =15.0 (C²⁸), 17.1 (C¹⁸), 17.7 (C²⁷), 20.8 [O-C(O)CH₃], 21.0 (C⁶), 22.5 (C³⁰), 22.6 (C²), 25.6 (C²⁶), 28.4 (C²³), 28.7 (C²²), 29.3 (C¹⁹), 29.5 (C²), 29.5 (C⁷), 30.0 (C¹), 33.2 (C⁴), 36.3 (C¹²), 36.4 (C¹⁰), 38.9 (C¹⁵), 39.5 (C¹), 39.6 (C⁸), 42.7 (C⁵), 43.7 (C¹³), 48.4 (C¹⁴), 50.6 (C⁹), 51.3 (COOCH₃), 62.2 (C³), 66.9 (C³), 67.4 (C¹¹), 74.2 (C¹⁶), 123.0

(C²⁴), 128.2 (C^{5'}, C^{7'}), 130.1 (C²⁰), 132.3 (C²⁵), 145.0 (C^{6'}), 145.9 (C^{4'}, C^{8'}), 148.0 (C¹⁷), 170.2 (C²¹), 170.7 [O-C(O)CH₃]. Calcd, %: C 65.83; H 8.42; Br 10.95; N 3.84; O 10.96. C₄₀H₆₁BrN₂O₅. Found, %: C 65.89; H 8.40; Br 10.92; N 3.85. *m/z* 751.3 [M+Na]⁺, calcd for [C₄₀H₆₁BrN₂O₅Na]⁺ 751.4.

1-(3-(((3S,4S,8S,10S,11R,14S,16S, Z)-16-acetoxy-11-hydroxy-17-(1-methoxy-6-methyl-1-oxohept-5-en-2-ylidene)-4,8,10,14-tetramethylhexadecahydro-1H-cyclopenta[a]phenanthren-3-yl)amino)propyl)-4-(dimethylamino)pyridin-1-ium bromide (12)

Yield 0.75 g (63 %), white powder, mp: 186-188°C, [α]_D¹⁹ -5° (*c* 1.11, MeOH). ¹H NMR (MeOD): δ=0.81 s (3H, H¹⁸), 1.03 d (3H, H²⁸, *J* 6.5 Hz), 1.12-1.22 m (1H, H^{15a}), 1.17 s (3H, H¹⁹), 1.18-1.28 m (1H, H^{7a}), 1.24 s (3H, H³⁰), 1.45-1.50 m (1H, H⁹), 1.51 s (3H, H²⁷), 1.51-1.65 m (1H, H^{7b}), 1.53-1.60 m (2H, H⁶), 1.55-1.60 m (1H, H^{2a}), 1.58 s (3H, H²⁶), 1.65-1.75 m (1H, H^{1a}), 1.71-1.79 m (1H, H⁴), 1.72-1.80 m (1H, H^{12a}), 1.74-1.80 m (1H, H^{2b}), 1.83-1.97 m (2H, H²), 1.85-1.95 m (1H, H^{1a}), 1.90 s [3H, O-C(O)CH₃], 1.94-2.14 m (2H, H²³), 2.01-2.14 m (1H, H⁵), 2.03-2.14 m (1H, H^{1b}), 2.13-2.25 m (1H, H^{15b}), 2.20-2.29 m (1H, H^{12b}), 2.27-2.45 m (2H, H²²), 2.34-2.58 m (1H, H^{1b}), 2.96 d (1H, H¹³, *J* 11.2 Hz), 3.02-3.14 m (1H, H³), 3.09 s (6H, H⁹, H¹⁰), 3.18-3.56 m (2H, H³), 3.56 s (3H, COOCH₃), 4.20-4.33 m (1H, H¹¹), 5.01 t (1H, H²⁴, *J* 6.5 Hz), 5.73 d (1H, H¹⁶, *J* 8.1 Hz), 6.63 d (2H, H⁵, H⁷, *J* 5.6 Hz), 8.10 d (2H, H⁴, H⁸, *J* 5.6 Hz). ¹³C NMR (MeOD): δ=14.3 (C²⁸), 17.1 (C¹⁸), 17.7 (C²⁷), 18.7 (C⁶), 20.9 [O-C(O)CH₃], 22.3 (C³⁰, C²), 25.6 (C²⁶), 28.4 (C²³), 28.6 (C¹⁹), 28.8 (C²²), 29.6 (C⁷), 29.6 (C²), 30.9 (C⁴), 31.7 (C¹), 36.3 (C¹²), 36.6 (C¹⁰), 38.9 (C¹⁵), 39.7 (C⁹, C¹⁰), 39.8 (C⁸), 42.0 (C⁵), 43.9 (C¹³), 48.5 (C¹), 49.1 (C¹⁴), 50.4 (C⁹), 51.3 (COOCH₃), 63.7 (C³), 66.8 (C¹¹), 67.5 (C³), 74.3 (C¹⁶), 106.7 (C^{5'}, C^{7'}), 123.0 (C²⁴), 130.1 (C²⁰), 133.1 (C²⁵), 143.0 (C^{4'}, C^{8'}), 148.0 (C¹⁷), 156.2 (C^{6'}), 170.5 (C²¹), 170.8 [O-C(O)CH₃]. Calcd, %: C 65.27; H 8.61; Br 10.34; N 5.44; O 10.35. C₄₂H₆₆BrN₃O₅. Found, %: C 65.34; H 8.59; Br 10.31; N 5.45. *m/z* 794.5 [M+Na]⁺, calcd for [C₄₂H₆₆BrN₃O₅Na]⁺ 794.4.

1-(3-(((3S,4S,8S,10S,11R,14S,16S, Z)-16-acetoxy-11-hydroxy-17-(1-methoxy-6-methyl-1-oxohept-5-en-2-ylidene)-4,8,10,14-tetramethylhexadecahydro-1H-cyclopenta[a]phenanthren-3-yl)amino)propyl)-1-methylpiperidin-1-ium bromide (13)

Yield 0.75 g (65 %), white powder, mp: 169-171°C, [α]_D¹⁹ +6° (*c* 1.10, MeOH). ¹H NMR (MeOD): δ=0.85 s (3H, H¹⁸), 1.06 d (3H, H²⁸, *J* 6.8 Hz), 1.08 s (3H, H¹⁹), 1.14-1.25 m (1H, H^{15a}), 1.18-1.26 m (1H, H^{7a}), 1.30 s (3H, H³⁰), 1.50-1.57 m (1H, H⁹), 1.52-1.58 m (2H, H⁶), 1.55 s (3H, H²⁷), 1.56-1.63 m (2H, H^{2a}, H^{7b}), 1.58-1.70 m (4H, H⁵, H⁷), 1.63-1.70 m (1H, H^{1a}), 1.65 s (3H, H²⁶), 1.72-1.83 m (1H, H⁴), 1.75-1.89 m (2H, H^{2b}, H^{12a}, 2H, H²), 1.82-1.92 m (2H, H⁶), 1.88-1.96 m (1H, H^{1a}), 1.94 s [3H, O-C(O)CH₃], 1.94-2.12 m (2H, H²³), 2.00-2.09 m (2H, H^{1b}, H⁵), 2.10-2.21 m (1H, H^{15b}), 2.19-2.28 m (1H, H^{12b}), 2.31-2.48 m (2H, H²²), 2.59-2.66 m (1H, H^{1b}), 2.95-3.08 m (4H, H⁴, H⁸), 2.97-3.10 m (1H, H¹³), 3.04-3.12 m (1H, H³), 3.40 s (3H, H⁹), 3.60 s (3H, COOCH₃), 4.09-4.17 m (2H, H³), 4.25-4.34 m (1H, H¹¹), 5.04 t (1H, H²⁴, *J* 6.2 Hz), 5.70 d (1H, H¹⁶, *J* 8.1 Hz). ¹³C NMR (MeOD): δ=14.4 (C²⁸), 17.1 (C¹⁸), 17.9 (C²⁷), 18.6 (C⁶), 20.9 [O-C(O)CH₃], 22.1 (C²), 22.3 (C^{5'}, C^{7'}), 22.3 (C^{6'}), 22.6 (C³⁰), 25.6 (C²⁶), 28.3 (C²³), 28.6 (C¹⁹), 28.8 (C²²), 29.6 (C², C⁷), 30.9 (C⁴), 31.7 (C¹), 36.3 (C¹²), 36.7 (C¹⁰), 38.9 (C¹⁵), 39.9 (C⁸), 42.1 (C⁵), 43.8 (C¹³), 44.5 (C^{4'}, C^{8'}), 48.5 (C¹⁴), 48.7 (C¹), 50.0 (C⁹), 50.5 (C⁹), 51.4 (COOCH₃), 65.6 (C³), 66.7 (C¹¹), 67.6 (C³), 74.4 (C¹⁶), 122.9 (C²⁴), 130.3 (C²⁰), 132.6 (C²⁵), 147.8 (C¹⁷), 170.6 (C²¹), 170.9 [O-C(O)CH₃]. Calcd, %: C 65.67; H 9.27; Br 10.66; N 3.74; O 10.67. C₄₁H₆₉BrN₂O₅. Found, %: C 65.61; H 9.29; Br 10.68; N 3.73. *m/z* 771.4 [M+Na]⁺, calcd for [C₄₁H₆₉BrN₂O₅Na]⁺ 771.4.

Results and Discussion

The quaternization reaction is one of the methods for functionalizing organic molecules, involving the introduction of a quaternary ammonium group into the chemical structure. Several variants of this reaction exist, but the most common method entails alkylation of tertiary amines with alkyl halides [36]. Bromination of the carboxylic group of FA was carried out by reacting compound **1** with 1,2-dibromoethane under reflux in dry acetone in the presence of K₂CO₃ as base. The reaction afforded bromide **2** with a yield of 97 % (Fig. 1) [35]. Bromide **2** was then subjected to quaternization with tertiary aliphatic and aromatic amines, such as triethylamine, pyridine, N, N-dimethylaminopyridine, and N-methylpiperidine. The reaction proceeded in DMF at 70 °C. After workup and purification by column chromatography, quaternized derivatives **3–6** were isolated in 62–70 % yields (Fig. 1). The lack of a pronounced dependence of the yields on the nucleophilicity of the amine can be attributed to the dominant role of steric hindrance from the fusidane scaffold and the use of a large excess of the amine, which ensures high conversion regardless of the nucleophile's electronic properties.






Figure 1. Synthesis of compounds 3-6

The transformations of compound **2** were confirmed by the **NMR** spectroscopy of the products, particularly by the downfield shift of the aliphatic chain carbon signal of the ester fragment attached to the quaternary nitrogen (δ_c 55.66–60.23 ppm, depending on the substituent at the quaternary nitrogen) compared to the starting bromide (δ_c = 28.35 ppm). Moreover, in the ^1H NMR spectra of compounds **4** and **5**, aromatic proton signals appeared at δ 6.88–9.14 ppm, while for derivative **3**, the ^{13}C NMR spectrum showed additional signals for the terminal methyl groups of the triethylammonium fragment at δ = 8.20 ppm.

The reductive amination of aldehydes and ketones is one of the most important methods in the pharmaceutical industry due to its versatility and selectivity, enabling the one-step synthesis of amines with varying degrees of substitution [37]. Using reductive amination, amine **9** was obtained from the 3,11-dioxo methyl ester derivative of FA (**8**) by reacting the latter with 3-bromopropylamine in MeOH in the presence of $\text{Ti}(\text{O}i\text{-Pr})_4$ catalyst, followed by reduction of the reaction mixture with NaBH_4 (Fig. 2). The reaction proceeded selectively at the 3-keto group, while the 11-keto group did not undergo amination and was reduced to the hydroxyl function. The transformation of **8** into **9** was confirmed by NMR spectroscopy. In the ^{13}C NMR spectrum, signals of the two keto groups at δ_c = 215.95 and 209.55 ppm disappeared, and signals of the C atoms at newly formed N-CH-3 and O-CH-11 bonds appeared at δ = 67.42 and 66.98 ppm, respectively. The signal of carbon atom attached to bromine was registered at δ = 38.73 ppm. In the ^1H NMR spectrum of **9**, signals of the CH-3 protons (δ = 3.04–3.15 ppm) and CH-11 protons (δ = 4.22–4.35 ppm) were observed, along with a series of multiplets (δ = 1.14–1.27, 1.87–2.66, and 2.66–3.38 ppm) corresponding to the CH_2 groups of the bromoaminopropane fragment. NOESY spectra confirmed that C-3 acquired β -configuration upon reductive amination, while C-11 retained the α -configuration of the native molecule.

Figure 2. Synthesis of compound **9**

Compound **9** was involved into the reaction with triethylamine, pyridine, N, N-dimethylaminopyridine, and N-methylpiperidine in DMF upon heating, affording amphiphilic derivatives **10–13** in 60–67 % yields (Fig. 3). A similar trend was observed for this series, where the yields remained largely unaffected by the amine's nucleophilicity, further supporting the steric control exerted by the bulky triterpenoid core. The structures of the obtained compounds were confirmed by a comprehensive set of spectral studies, including 1D and 2D NMR spectroscopy data.

Figure 3. Synthesis of quaternized derivatives **10–13** from compound **9**

Conclusions

Brominated analogs of fusidane triterpenoids were synthesized via esterification of the FA carboxylic group with 1,2-dibromoethane and by reductive amination of the methyl fusidate diketone derivative with 3-bromopropylamine. Quaternization of aliphatic and aromatic tertiary amines with the resulting fusidane bromo derivatives afforded novel amphiphilic fusidic acid analogs, which are of interest as potential antibacterial agents with membrane-tropic activity.

Funding

This study was carried out in the framework of state assignment to the Institute of Petrochemistry and Catalysis (project no. FMRS-2025-0033).

Author Information*

*The authors' names are presented in the following order: First Name, Middle Name and Last Name

Elena Viktorovna Salimova — Candidate of Chemical Sciences, Research Fellow, Organic Synthesis Laboratory, Institute of Petrochemistry and Catalysis, Ufa Federal Research Center of the Russian Academy of Sciences, prosp. Oktyabrya, 141, 450075, Ufa, Russia; e-mail: salimovaev@mail.ru; <https://orcid.org/0000-0002-4328-5080>

Rinat Ravilevich Gubaidullin — Candidate of Chemical Sciences, Research Fellow, Organic Synthesis Laboratory, Institute of Petrochemistry and Catalysis, Ufa Federal Research Center of the Russian Academy of Sciences, prosp. Oktyabrya, 141, 450075, Ufa, Russia; e-mail: rinatg83@mail.ru; <https://orcid.org/0000-0002-9915-0820>

Lyudmila Vyacheslavovna Parfenova (*corresponding author*) — Doctor of Chemical Sciences, Head of Organic Synthesis Laboratory, Institute of Petrochemistry and Catalysis, Ufa Federal Research Center of the Russian Academy of Sciences, prosp. Oktyabrya, 141, 450075, Ufa, Russia; e-mail: lu-da_parfenova@mail.ru; <https://orcid.org/0000-0003-2816-2178>

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript. **CRedit**: **Elena Viktorovna Salimova** – data curation, investigation, methodology, validation, visualization, writing—original draft; **Rinat Ravilevich Gubaidullin** – formal analysis, writing—original draft; **Lyudmila Vyacheslavovna Parfenova** – conceptualization, funding acquisition, resources, supervision, validation, writing-review & editing.

Acknowledgments

Structural studies of the synthesized compounds were performed with the use of the Collective Usage Centre “Agidel” at the Institute of Petrochemistry and Catalysis of the Russian Academy of Sciences (RAS).

Conflicts of Interest

The authors declare no conflict of interest.

References

- Gerba, C. P. (2015). Quaternary ammonium biocides: Efficacy in application. *Applied and Environmental Microbiology*, 81(2), 464–469. <https://doi.org/10.1128/AEM.02633-14>
- Chanawanno, K., Thuptimandang, P., Chantrapromma, S., & Fun, H. K. (2022). New tunable pyridinium benzenesulfonate amphiphiles as anti-MRSA quaternary ammonium compounds (QACs) *Journal of Molecular Structure*, 1254, 132389. <https://doi.org/10.1016/j.molstruc.2022.132389>
- Walker, E. B. (2002). Chapter 5. In D.S. Paulson (Ed.), *Handbook of topical antimicrobials: Industrial applications in consumer products and pharmaceuticals*. CRC Press. <https://doi.org/10.1201/9780203909256>
- Savage, P. B., Li, C., Taotafa, U., Ding, B., & Guan, Q. (2002). Antibacterial properties of cationic steroid antibiotics. *FEMS Microbiology Letters*, 217(1), 1–7. <https://doi.org/10.1111/j.1574-6968.2002.tb11448.x>
- Wessels, S., & Ingmer, H. (2013). Modes of action of three disinfectant active substances: a review. *Regulatory Toxicology and Pharmacology*, 67, 456–467. <https://doi.org/10.1016/j.yrtph.2013.09.006>
- Jennings, M. C., Minbiole, K. P. C., & Wuest, W. M. (2015). Quaternary Ammonium Compounds: An Antimicrobial Mainstay and Platform for Innovation to Address Bacterial Resistance. *ACS Infectious Diseases*, 1, 288–303. <https://doi.org/10.1021/acsinfecdis.5b00047>
- Schrank, C. L., Minbiole, K. P. C., & Wuest, W. M. (2020). Are Quaternary Ammonium Compounds, the Workhorse Disinfectants, Effective against Severe Acute Respiratory Syndrome-Coronavirus-2?. *ACS Infectious Diseases*, 6, 1553–1557. <https://doi.org/10.1021/acsinfecdis.0c00265>
- Chanawanno, K., Chantrapromma, S., Anantapong, T., Kanjana-Opas, A., & Fun, H. K. (2010). Synthesis, structure and *in vitro* antibacterial activities of new hybrid disinfectants quaternary ammonium compounds: Pyridinium and quinolinium stilbene benzenesulfonates. *European Journal of Medicinal Chemistry*, 45, 4199–4208. <https://doi.org/10.1016/j.ejmech.2010.06.014>
- Forman, M. E., Jennings, M. C., Wuest, W. M., & Minbiole, K. P. C. (2016). Building a Better Quaternary Ammonium Compound (QAC): Branched Tetracationic Antiseptic Amphiphiles. *ChemMedChem*, 11, 1401–1405. <https://doi.org/10.1002/cmdc.201600176>

- 10 Joyce, M. D., Jennings, M. C., Santiago, C. N., Fletcher, M. H., Wuest, W. M., & Mibiole, K. P. (2016). Natural product-derived quaternary ammonium compounds with potent antimicrobial activity. *Journal of Antibiotics*, 69, 344–347. <https://doi.org/10.1038/ja.2015.107>
- 11 Sapozhnikov, S. V., Sabirova, A. E., Shtyrlin, N. V., Druk, A. Y., Agafonova, M. N., Chirkova, M. N., Kazakova, R. R., Grishaev, D. Y., Nikishova, T. V., Krylova, E. S., Nikitina, E. V., Kayumov, A. R., & Shtyrlin, Y. G. (2021). Design, synthesis, antibacterial activity and toxicity of novel quaternary ammonium compounds based on pyridoxine and fatty acids. *European Journal of Medicinal Chemistry*, 211, 113100. <https://doi.org/10.1016/j.ejmech.2020.113100>
- 12 Yamanaka, T., Bannai, H., Tsujimura, K., Nemoto, M., Kondo, T., & Matsumura, T. (2014). Comparison of the Virucidal Effects of Disinfectant Agents Against Equine Influenza A Virus. *Journal of Equine Veterinary Science*, 34, 715–718. <https://doi.org/10.1016/j.jevs.2013.12.010>
- 13 Nyström, L., & Malmsten, M. (2018). Membrane interactions and cell selectivity of amphiphilic anticancer peptides. *Current Opinion in Colloid & Interface Science*, 38, 1–17. <https://doi.org/10.1016/j.cocis.2018.06.009>
- 14 Yang, J. S., Song, D., Lee, B., Ko, W. J., Park, S. K., Won, M., Lee, K., Kim, H. M., & Han, G. (2011). Synthesis and biological evaluation of novel aliphatic amido-quaternary ammonium salts for anticancer chemotherapy: Part I. *European Journal of Medicinal Chemistry*, 46, 2861–2866. <https://doi.org/10.1016/j.ejmech.2011.04.009>
- 15 Song, D., Yang, J. S., Oh, C., Cui, S., Kim, B. K., Won, M., Lee, J. I., Kim, H. M., & Han, G. (2013). New synthetic aliphatic sulfonamido-quaternary ammonium salts as anticancer chemotherapeutic agents. *European Journal of Medicinal Chemistry*, 69, 670–677. <https://doi.org/10.1016/j.ejmech.2013.09.022>
- 16 Dubinin, M. V., Semenova, A. A., Ilzorkina, A. I., Penkov, N. V., Nedopekina, D. A., Sharapov, V. A., Khoroshavina, E. I., Davletshin, E. V., Belosludtseva, N. V., Spivak, A. Y., Belosludtsev, K. N., et al. (2021). Mitochondria-targeted prooxidant effects of betulinic acid conjugated with delocalized lipophilic cation F16. *Free Radical Biology and Medicine*, 168, 55–69. <https://doi.org/10.1016/j.freeradbiomed.2021.03.036>
- 17 Dey, G., Bharti, R., Sen, R., & Mandal, M. (2015). Microbial amphiphiles: a class of promising new-generation anticancer agents. *Drug Discovery Today*, 20, 136–146. <https://doi.org/10.1016/j.drudis.2014.09.006>
- 18 Ciunac, D., Gong, H., Hu, X., & Lu, J. R. (2019). Membrane targeting cationic antimicrobial peptides. *Journal of Colloid and Interface Science*, 537, 163–185. <https://doi.org/10.1016/j.jcis.2018.10.103>
- 19 Dubinin, M. V., Semenova, A. A., Ilzorkina, A. I., Markelova, N. Y., Penkov, N. V., Shakurova, E. R., Belosludtsev, K. N., & Parfenova, L. V. (2021). New quaternized pyridinium derivatives of betulin: Synthesis and evaluation of membranotropic properties on liposomes, pro- and eukaryotic cells, and isolated mitochondria. *Chemico-Biological Interactions*, 349, 109678. <https://doi.org/10.1016/j.cbi.2021.109678>
- 20 Pasquale, E. D., Salmi-Smail, C., Brunel, J. M., Sanchez, P., Fantini, J., & Maresca, M. (2010). Biophysical studies of the interaction of squalamine and other cationic amphiphilic molecules with bacterial and eukaryotic membranes: Importance of the distribution coefficient in membrane selectivity. *Chemistry and Physics of Lipids*, 163, 131–140. <https://doi.org/10.1016/j.chemphyslip.2009.10.006>
- 21 Krasutsky, P. A., Avilov, D., Sergeeva, T., Krasutsky, D. A., & Kolomitsyna, O. (2008). *US Patent No. 7,199,114 B2*.
- 22 Kataev, V. E., Strobkyina, I. Yu., & Zakharova, L. Ya. (2014). Quaternary ammonium derivatives of natural terpenoids. Synthesis and properties. *Russian Chemical Bulletin*, 63, 1884–1900. <https://doi.org/10.1007/s11172-014-0680-x>
- 23 Zhao, M., Godecke, T., Gunn, J., Duan, J. A., & Che, C. T. (2013). Protostane and Fusidane Triterpenes: A Mini-Review. *Molecules*, 18, 4054–4080. <https://doi.org/10.3390/molecules18044054>
- 24 Dziwornu, G. A., Kamunya, S., Ntsabo, T., & Chibale, K. (2019). Novel antimycobacterial C-21 amide derivatives of the antibiotic fusidic acid: synthesis, pharmacological evaluation and rationalization of media-dependent activity using molecular docking studies in the binding site of human serum albumin. *MedChemComm*, 10, 961–969. <https://doi.org/10.1039/c9md00161a>
- 25 Guo, M., Ren, Q., Wang, B., Ji, W., Ni, J., Feng, Y., Bi, Y., Tian, J., & Wang, H. (2019). Discovery and synthesis of 3- and 21-substituted fusidic acid derivatives as reversal agents of P-glycoprotein-mediated multidrug resistance. *European Journal of Medicinal Chemistry*, 182, 111668. <https://doi.org/10.1016/j.ejmech.2019.111668>
- 26 Ni, J., Guo, M., Cao, Y., Lei, L., Liu, K., Wang, B., Lu, F., Zhai, R., Gao, X., Yan, C., Wang, H., & Bi, Y. (2019). Discovery, synthesis of novel fusidic acid derivatives possessed amino-terminal groups at the 3-hydroxyl position with anticancer activity. *European Journal of Medicinal Chemistry*, 162, 122–131. <https://doi.org/10.1016/j.ejmech.2018.10.059>
- 27 Espinoza-Moraga, M., Singh, K., Njoroge, M., Kaur, G., Okombo, J., De Kock, C., Smith, P. J., Wittlin, S., & Chibale, K. (2017). Synthesis and biological characterisation of ester and amide derivatives of fusidic acid as antiplasmodial agents. *Bioorganic & Medicinal Chemistry Letters*, 27, 658–661. <https://doi.org/10.1016/j.bmcl.2016.11.077>
- 28 Salimova, E. V., Magafurova, A. A., Tretyakova, E. V., Kukovinets, O. S., & Parfenova, L. V. (2020). Indole Derivatives of Fusidane Triterpenoids: Synthesis and the Antibacterial Activity. *Chemistry of Heterocyclic Compounds*, 56, 800–804. <https://doi.org/10.1007/s10593-020-02733-1>
- 29 Salimova, E. V., & Parfenova, L. V. (2021). Synthesis and Biological Activity of Oximes, Amines, and Lactams of Fusidane Triterpenoids. *ChemistrySelect*, 6, 8848–8854. <https://doi.org/10.1002/slct.202102019>
- 30 Salimova, E. V., Mamaev, A. G., Tretyakova, E. V., Kukovinets, O. S., Mavzyutov, A. R., Shvets, K. Yu., & Parfenova, L. V. (2018). Synthesis and Biological Activity of Cyanoethyl Derivatives of Fusidic Acid. *Russian Journal of Organic Chemistry*, 54, 1411–1418. <https://doi.org/10.1134/S1070428018090245>
- 31 Salimova, E. V., Mamaev, A. G., Tretyakova, E. V., Kukovinets, O. S., & Parfenova, L. V. (2018). Reductive amination of fusidane triterpenoid ketones. *Mediterranean Journal of Chemistry*, 7, 198–203. <https://doi.org/10.13171/mjc7318106921-salimova>

- 32 Salimova, E. V., Tret'yakova, E. V., & Parfenova, L. V. (2019). Synthesis and cytotoxic activity of 3-amino substituted fusidane triterpenoids. *Medicinal Chemistry Research*, 28, 2171–2183. <https://doi.org/10.1007/s00044-019-02445-y>
- 33 Salimova, E. V., Tret'yakova, E. V., Kukovinets, O. S., & Parfenova, L. V. (2020). Beckmann Rearrangement of Oximes of the Fusidane Series. *Russian Journal of Organic Chemistry*, 56, 11–19. <https://doi.org/10.31857/S0514749220010036>
- 34 Salimova, E. V., & Parfenova, L. V. (2022). Fischer Reaction in the Synthesis of New Triterpene Indoles of the Fusidane Series. *Russian Journal of Organic Chemistry*, 58, 25–37. <https://doi.org/10.1134/S1070428022010031>
- 35 Salimova, E. V., Mannanova, L. R., Kukovinets, O. S., & Parfenova, L. V. (2022). Synthesis of Halo Derivatives of Fusidane Triterpenoids. *Russian Journal of Organic Chemistry*, 58, 968–976. <https://doi.org/10.1134/S1070428022070041>
- 36 Geresh, S., Dawadi, R. P., & Arad, S. M. (2000). Chemical modifications of biopolymers: quaternization of the extracellular polysaccharide of the red microalga *Porphyridium* sp. *Carbohydrate Polymers*, 43, 75–80. [https://doi.org/10.1016/S0144-8617\(99\)00194-0](https://doi.org/10.1016/S0144-8617(99)00194-0)
- 37 Vila, C., Díaz-García, P., Ladeira, S., Balerdi, M., & Freixa, Z. (2020). Catalytic reductive amination of aldehydes and ketones with nitro compounds: New light on an old reaction. *Frontiers in Chemistry*, 8, 215. <https://doi.org/10.3389/fchem.2020.00215>