

SUPPLEMENTARY MATERIALS

Development of Novel Wheat Waste-Derived Biochar and Its Potential in Pharmaceutical Wastewater Treatment

Ayazhan M. Turarbek¹, Farida Zh. Abilkanova², Aitolkyn S. Uali^{1*}

¹ *L.N. Gumilyov Eurasian National University, Astana, Kazakhstan*

² *Karaganda Industrial University, Temirtau, Kazakhstan*

(*Corresponding author's e-mail: uali_as@enu.kz)

List of Content

FTIR spectrum of the intermediate biochar product after the first carbonization stage (Figure S1).	S2
FTIR peak assignments to the FTIR spectrum given in Figure S1 (Table S1).	S3
References	S4



Figure S1. FTIR spectrum of the intermediate biochar product after the first carbonisation stage

Table S1

FTIR peak assignments to the FTIR spectrum given in Figure S1

FTIR peak position	Functional group(s)	Ref-s
3295	O-H stretch (carboxylic group or alcohol)	[1]
2906, 2875	C-H stretch (alkyl CH)	[1,2]
1599, 1552	C=C stretch (aryl, double-bond)	[2]
1399, 1373	CH ₃ -, CH ₂ - (aliphatic chains)	[3]
1312	C-H stretch (aliphatic chains)	[4]
≈1141	O-H stretch (carboxylic group) Aromatic ring stretching vibration or C-O stretching vibration	[5] [3]
1024	C-OC/C-OH stretches	[1]
941 (995-905)	C-O stretch in cellulose	[6]
≈595-593	Out-of-plane-vibrations of associated OH groups	[7]

References

- 1 Adeniyi, A. G., Adeyanju, C. A., Emenike, E. C., Otoikhian, S. K., Ogunniyi, S., Iwuozor, K. O., & Raji, A. A. (2022). Thermal energy recovery and valorisation of *Delonix regia* stem for biochar production. *Environmental Challenges*, 9, 100630. <https://doi.org/10.1016/j.envc.2022.100630>
- 2 Chia, C. H., Gong, B., Joseph, S. D., Marjo, C. E., Munroe, P., & Rich, A. M. (2012). Imaging of mineral-enriched biochar by FTIR, Raman and SEM–EDX. *Vibrational Spectroscopy*, 62, 248–257. <https://doi.org/10.1016/j.vibspec.2012.06.006>
- 3 Guo, H., Ma, L., Shen, F., Yang, G., Zhang, Y., Deng, S., Zhang, J., Song, C., & Zeng, Y. (2017). Effects of La-involvement on biomass pyrolysis behaviors and properties of produced biochar. *Journal of Rare Earths*, 35(6), 593–601. [https://doi.org/10.1016/s1002-0721\(17\)60952-9](https://doi.org/10.1016/s1002-0721(17)60952-9)
- 4 Zhao, J.-J., Huang, X.-H., Hua, L.-L., Zhou, S.-S., Jiang, W., Tang, Y.-C., & Qian, J. (2022). Effects of Tea Residue Biochar on Phosphorus Adsorption-Desorption in Soil. *Polish Journal of Environmental Studies*, 31(3), 2461–2471. <https://doi.org/10.15244/pjoes/143357>
- 5 Liang, H., Zhu, C., Ji, S., Kannan, P., & Chen, F. (2022). Magnetic Fe₂O₃/biochar composite prepared in a molten salt medium for antibiotic removal in water. *Biochar*, 4(1). <https://doi.org/10.1007/s42773-021-00130-1>
- 6 Reza, M. S., Afroze, S., Bakar, M. S. A., Saidur, R., Aslfattahi, N., Taweekun, J., & Azad, A. K. (2020). Biochar characterization of invasive *Pennisetum purpureum* grass: effect of pyrolysis temperature. *Biochar*, 2(2), 239–251. <https://doi.org/10.1007/s42773-020-00048-0>
- 7 Ilić, M., Haegel, F.-H., Lolić, A., Nedić, Z., Tosti, T., Ignjatović, I. S., Linden, A., Jablonowski, N. D., & Hartmann, H. (2022). Surface functional groups and degree of carbonization of selected chars from different processes and feedstock. *PLOS ONE*, 17(11), e0277365. <https://doi.org/10.1371/journal.pone.0277365>